arXiv:1603.01087v1 [physics.atom-ph] 3 Mar 2016

Compilation of spectroscopic data of Radium (Ra I and Ra II)

U. Dammalapati,[1, a)] K. Jungmann,[1] and L. Willmann[1, b)]
Van Swinderen Institute, Faculty of Mathematics and Natural Sciences, University of Groningen, Zernikelaan 25, 9747 AA Groningen, The Netherlands

(Dated: 4 March 2016)

Energy levels, wavelengths, lifetimes and hyperfine structure constants for the isotopes of the first and second spectra of radium, Ra I and Ra II have been compiled. Wavelengths and wave numbers are tabulated for ^{226}Ra and for other Ra isotopes. Isotope shifts and hyperfine structure constants of even and odd-A isotopes of neutral radium atom and singly ionized radium are included. Experimental lifetimes of the states for both neutral and ionic Ra are also added, where available. The information is beneficial for present and future experiments aimed at different physics motivations using neutral Ra and singly ionized Ra.

Keywords: Radium (Ra I); Radium ion (Ra II); energy levels; wave numbers; transitions; spectroscopic data; hyperfine structure; lifetime.

I. INTRODUCTION

Radium (Ra), a radioactive element was discovered by P. Curie, M. Curie and G. Bemont in the year 1898[1]. It was extracted by chemical means from pitchblende or uraninite. About one gram (1g) of radium is present in 7 tons of pitchblende. The activity of 1g of ^{226}Ra was defined later as the unit of radioactivity, Curie, Ci (1 Ci = 3.7×10^{10} disintegrations per second (dis/s)). The term *radioactivity* was introduced by Mme. Marie Curie. Ritz in 1908 classified two series of lines of neutral radium, Ra I[2]. In 1911, radium was isolated by Mme. Curie and Debierne, who found it to be a brilliant white solid[3]. Some energy levels and line classifications of Ra I were given by Fues in 1920[4]. In 1922, Fowler gave classifications of some series of lines in both Ra I and singly ionized radium, Ra II[5]. In 1933-34, Rasmussen first measured the energy levels of singly ionized radium (Ra II)[6] using arc discharge and of neutral radium (Ra I)[7] using 2 mg of radiumchloride ($RaCl_2$) in a discharge lamp and monochromator for determining the wavelengths. Some of the transition terms of neutral radium measured by Rasmussen were corrected by H. N. Russell[8]. The first compilation of energy levels for neutral and singly ionized radium was published by Moore in 1958 in *Atomic Energy Levels* (AEL)[9]. After that, no updates were made to reference data on neutral and singly ionized Ra. With the advent of accelerator facilities, modern experimental methods to produce radium isotopes and to probe the atomic structure, more information is available on these spectra in the literature. At this juncture, we realized the need to update the information available on Ra atom and Ra ion and gathered it, which might be beneficial for various experiments that are in progress[10–14] and for the future experiments[15,16] with different physics motivations and for various fields of research.

[a)] dammalapati.umakanth.a8@tohoku.ac.jp; Present address: Cyclotron and Radioisotope Center (CYRIC), Tohoku University, 6-3 Aramaki-aza Aoba, Aoba-ku, Miyagi 980-8578, Japan
[b)] l.willmann@rug.nl

About forty six years after Rasmussen's measurement of energy levels and wavelengths in Ra ion and Ra atom, furnace-absorption measurements of energy levels of the $7snp\ ^1P_1^0$ ($n = 13 - 52$) series of neutral radium in the ultraviolet region were reported by Armstrong et al.[17,18]. From the experimental results and applying the multichannel quantum defect theory (MQDT), values for $7snp$ ($n = 9-12$) levels were semi-empirically determined. The reported uncertainty was 0.006 cm^{-1} for wavenumbers and 0.001 $\mathring{A}$ for wavelengths.

The first systematic measurements on a series of neutral and singly ionized radium isotopes employing the technique of collinear laser spectroscopy were conducted by the ISOLDE collaboration at CERN, Geneva[19–23]. Radium isotopes were produced by impinging protons of energy 600 MeV on a UC_2 cloth target. These studies established nuclear spins and obtained nuclear moments from the measured hyperfine structure and isotope shifts.

In the last decade, there has been renewed interest in radium element (radium atom as well as ion) from both theoretical and experimental physicists. From theoretical calculations, radium atom is proposed as a promising candidate for searches of a permanent electric dipole moment in an atomic system[24–30]. The reasons are: large polarizability due to the near degeneracy of the opposite parity $7s7p\ ^3P_1^0$ and $7s6d\ ^3D_2$ levels; some isotopes of radium have nuclear octupole deformation; and high atomic number ($Z = 88$). Singly ionized radium ion is proposed as a promising candidate for atomic parity violation experiments and as an atomic clock[31–33].

In this context, short lived radioactive isotopes were produced at the TRIμP facility, University of Groningen, The Netherlands[34]. Radium isotopes were produced by the method of inverse reaction kinematics using ^{206,204}Pb beams hitting a rotating carbon target[35,36]. Precision laser spectroscopy of radium ions was performed in a radio frequency quadrupole (RFQ) trap[13,37–39].

At the University of Groningen, The Netherlands, with ^{229}Th (10 μCi) placed in an oven, ^{225}Ra neutral atomic beam was produced. Employing this thermal beam of Ra atoms, the absolute frequency of the strong electric dipole $7s^2\ \ ^1S_0$ - $7s7p\ ^1P_1^0$ transition and weaker in-

tercombination $7s^2\ ^1S_0$ - $7s7p\ ^3P_1$ transition were reported with reference to a femtosecond optical frequency comb[12,40]. The uncertainty reported was 0.00006 cm^{-1} and 0.00013 cm^{-1}, respectively.

At Argonne National Laboratory (ANL), USA, electrochemically separated neutral radium isotopes, ^{225,226}Ra placed in an oven were used for a laser cooling and trapping experiment[10]. Further, optical dipole trapping of ^{226}Ra[11] and measurement of lifetimes of $7s7p\ ^3P_1^0$, $7s6d\ ^3D_1$ and $7s6d\ ^1D_2$ states were reported by the ANL group[10,41,42]. Very recently, using cold atoms, the first measurement of the atomic dipole moment of ^{225}Ra was reported by the same group[14].

In this compilation, energy levels, wavelengths, wave numbers, transition probabilities, lifetimes and other spectral data have been reviewed. The most accurate information has been assembled from the available experimental and theoretical data and is presented in Tables I - XVI. The uncertainty of each value, as given by the original authors, is indicated in the tables. In the line tables (Tables III and X), the wavelengths and their uncertainties are reported in units of angstroms. For lines between 2000 $\mathring{A}$ and 15000 $\mathring{A}$ air wavelengths and for lines below 2000 $\mathring{A}$ vacuum wavelengths are given. The index of refraction is determined by the three-term formula (Equation 3) of Peck and Reeder[43]. The wave number of each transition is given in reciprocal centimeters and its uncertainty is also given wherever applicable. The lower and upper level columns indicate the classification given for the transition. The Ritz wavelengths and their uncertainties calculated using the level optimization code, LOPT[44] are also given. The calculated transition probabilities (A_{ik}) are given in units of inverse seconds. Their accuracy range is also given in a separate column. For singly ionized Ra uncertainty is given for theoretical transition rates by Sahoo *et al.*[32] (Table X).

The energy level tables (Tables II and IX) contain the configuration, term, and J values of each energy level, using LS coupling to describe the configurations. In addition to that, energy uncertainty, energy level values and their uncertainty from the LOPT code are also given. For visual clarity, only the first member of the term has the configuration written out similar to the notation used by Curry[45]. The level value and its uncertainty are given in reciprocal centimeters.

Ionization energies are given in units of cm^{-1} and electronvolts (eV). Conversion factor from cm^{-1} to eV was taken from the 2014 CODATA adjustment of fundamental constants[46]: 1 cm^{-1} = 1.239 841 9739(76) $\times\ 10^{-4}$ eV. The reciprocal centimeter is related to the SI unit for energy, the joule, in the following way: 1 cm^{-1} is equivalent to $1.986445824(24) \times 10^{-23}$ J[46].

The hyperfine structure constants (HFS) and isotope shifts are given in Tables V-VIII, XIII-XV in units of megahertz (MHz) with the uncertainty in the last digit given in parentheses following the value. All uncertainties given in this paper are on the level of one standard deviation. The hyperfine splitting constant, A, is the magnetic dipole coefficient, whereas B is the electric quadrupole coefficient. The references for energy levels and hyperfine structure splitting and isotope shifts are given to the source of the energy level value and hyperfine splitting constants, respectively.

II. RA I

In Table I, radium isotopes, atomic weights, half-lives, nuclear spins, nuclear magnetic moments and electric quadrupole moments are given. Up to now 35 radium isotopes have been discovered excluding the isomers of ^{203m,205m,207m,213m}Ra isotopes.

Table II presents the energy levels of ^{226}Ra. They are taken from Moore 1958 compilation[9] and other sources where improved experimental values are available. Moore's compilation is mainly based on the measurements of Rasmussen[6] and corrections made to them by Russell[8]. In addition to the compilation of Moore we have added the energy levels of ^{226}Ra in the UV region from the absorption measurements of Armstrong *et al.*[17]. However, the uncertainty $\pm$0.006 cm^{-1} given by Armstrong *et al.* for wavenumbers is not correct as the uncertainty 0.001 $\mathring{A}$ in their wavelengths corresponds to the wavenumber uncertainty of 0.018 cm^{-1}. This is brought to the notice of the authors by Alexander Kramida[49]. The updated uncertainties of the wavenumbers are given in Table II. We have not included the semi-empirical values given in Armstrong *et al.*[17] for $7s9p - 7s12p$ energy levels. Improved values of the $7s7p\ ^1P_1^0$ level from Santra *et al.*,[12] $7s7p\ ^3P_1^0$ level from Scielzo *et al.*,[41] and $7s6d\ ^3D_1$ level from Guest *et al.*,[10] are also given. Furthermore, the measurements of Rasmussen and Armstrong *et* al. are attributed to the ^{226}Ra isotope due to its natural abundance and long half-life compared to the other Ra isotopes. Also, the high probability that Rasmussen used ^{226}Ra is supported by the fact that the knowledge of isotopes was not available at that time as the first isotope of an element observed and reported by Urey *et al.* in 1932 was deuterium[50]. Furthermore, the recent measurements from the Argonne group and Groningen group support this in addition to the ISOLDE group. The very recent measurement of the $7s6d\ ^3D_1$-$7s7p\ ^1P_1^0$ transition in ^{226}Ra by the Argonne group[10] confirms attribution of Rasmussen's measurements to the ^{226}Ra isotope.

The $7p^2$ configuration of the $^1D_2$ term given in Moore's compilation from Rasmussen measurements was changed to the $7s7d$ configuration by the ISOLDE group[21] based on the analysis of their experimental results. In Table II of neutral Ra energy level data it is shown in parentheses. For the $^1D_2$ term, the $6d^2$ configuration is attributed by Quinet *et al.*[51]. It is also shown in parentheses. Also, the $^3P_{0,1,2}$ levels assigned to the $7p^2$ configuration as given in Moore's compilation are predicted to belong to the $6d^2$ configuration by Quinet *et al.*[51], which is shown in parentheses. For the $7s8p\ ^1P_1^0$ level, Armstrong *et al.*[17] restored Rasmussen's identification. Furthermore, Gin-

TABLE I. Isotopes of radium with zero nuclear spin (I) and with non-zero nuclear spin. The half-lives (ns-nanoseconds, μs-microseconds, ms-milliseconds, s-seconds, m-minutes, d-days and y-years) and nuclear spin (I) are taken from Ref. [47]. The atomic weights are taken from Ref. [48]

Isotope	Atomic weight[48] u	Half-life $t_{1/2}$	Nuclear spin[19] I	Nuclear magnetic moment[47] $[\mu_I/\mu_N]$	Electric Quadrupole moment[23] Q_s in [b] [a]
^{235}Ra	235.054970(320)				
^{234}Ra	234.050340(30)	30 s	0		
^{233}Ra	233.047582(17)	30 s			
^{232}Ra	232.043475(10)	4.2 m	0		
^{231}Ra	231.041027(12)	104.1 s	5/2		
^{230}Ra	230.037055(11)	93 m	0		
^{229}Ra	229.034942(16)	4.0 m	5/2	0.5025(27)	3.09(4)
^{228}Ra	228.0310707(2.6)	5.75 y	0		
^{227}Ra	227.0291783(2.5)	42.2 m	3/2	-0.4038(24)	1.58(3)
^{226}Ra	226.0254103(2.5)	1600 y	0		
^{225}Ra	225.023612(3)	14.9 d	1/2	-0.7338(15)	
^{224}Ra	224.020212(2.3)	3.6319 d	0		
^{223}Ra	223.0185023(2.7)	11.43 d	3/2	0.2705(19)	1.254(3)
^{222}Ra	222.015375(5)	38.0 s	0		
^{221}Ra	221.013918(5)	28 s	5/2	-0.1799(17)	1.978(7)
^{220}Ra	220.011026(9)	18 ms	0		
^{219}Ra	219.010085(9)	10 ms	7/2		
^{218}Ra	218.007141(12)	25.2 μs	0		
^{217}Ra	217.006321(9)	1.6 μs	9/2		
^{216}Ra	216.003533(9)	182 ns	0		
^{215}Ra	215.002720(8)	1.55 ms	9/2		
^{214}Ra	214.000100(6)	2.46 s	0		
^{213}Ra	213.000384(22)	2.73 m	1/2	0.6133(18)	
^{212}Ra	211.999787(12)	13 s	0		
^{211}Ra	211.000893(9)	13 s	5/2	0.8780(38)	0.48(2)
^{210}Ra	210.000494(16)	3.7 s	0		
^{209}Ra	209.001990(50)	4.6 s	5/2	0.865(13)	0.40(2)
^{208}Ra	208.001841(17)	1.3 s	0		
^{207}Ra	207.003800(60)	1.35 s	3/2,5/2		
^{206}Ra	206.003828(19)	0.24 s	0		
^{205}Ra	205.006270(80)	210 ms	3/2		
^{204}Ra	204.006492(16)	57 ms	0		
^{203}Ra	203.009300(90)	31 ms	3/2		
^{202}Ra	202.009760(26)	16 ms	0		
^{201m}Ra	201.012710(110)	1.6 ms	13/2		

[a] 1 barn (b) = 10^{-24} cm^2

ges and Dzuba in their recent theoretical work[52] noted that their calculated value of the $7s8p\ ^1P_1$ energy level is very low compared to the values from Rasmussen's data and Moore's compilation. For example, the difference from Moore's compilation is 2155 cm^{-1} . Because of the discrepancies between the observed spectra and the contradicting interpretations of configuration terms of some energy levels and deviations from the theoretical calculations in Ra I more experimental data is required.

In Table III the intensity, observed wavelengths in air, uncertainty in observed wavelengths, observed wavenumbers, lower and upper levels, classifications, Ritz wavelengths obtained with the LOPT code, uncertainties in Ritz wavelengths and transition probabilities are given. The Ritz wavelengths are not given for lines that alone determine one of the levels involved in the transition. Some of the observed wavenumbers given in column 4 differ from the wavenumbers given in the original work of Rasmussen[7] and Armstrong *et al.*[17]. The difference from the original values and the present reported values is due to the conversion formulas used by Rasmussen and Armstrong *et al.* for arriving at the wavenumbers from the measured wavelengths. Here we have used the widely accepted formula of Peck and Reeder[43] to arrive at the wavenumbers. The first reference (Column 7) corresponds to the transitions/lines and the second reference (Column 11) to transition probabilities. We have given theoretical transition rates for some of the transitions in Ra atom calculated by Dzuba and Flambaum[53] and one value from the experiment[41]. In column 4 of Table III, we have given wavenumbers for the transitions from the measurements of Rasmussen[7] and from Russell[8]. Ab-

sorption measurements from Armstrong *et al.*[17] have no information on the intensity of the observed lines. Their wavenumbers and air wavelengths with uncertainties are also given. For the 7s^2 $^1S_0$-7s7p $^{3,1}P_1^0$ transitions two line references are given; the first one for the intensity and the second one for the wavenumber. The wavenumber and the air wavelength of the 7s6d $^3D_1$-7s7p $^3P_1^0$ transition measured by the ANL group is also given. The measured value is 6999.84 cm^{-1} with an uncertainty of 0.02 cm^{-1}. In columns 8 and 9, Ritz wavelengths and their uncertainties are given. These are obtained from the calculations using the LOPT code of National Institute of Standards and Technology (NIST), USA.

Theoretically calculated lifetimes and transition rates for some of the energy levels in neutral Ra have been previously reported[26,53,54]. However, only three experimental lifetimes of states in neutral Ra atom are available and they are given in Table IV.

In Table V, isotope shifts in frequency for the $7s^2$ ^{1}S$_0$-$7s7p$ ^{1}P$_1$ electric dipole allowed transition of 19 neutral Ra isotopes and hyperfine structure constants A and B of the $7s7p$ ^{1}P$_1$ level for the odd isotopes are given. The isotope shifts are given with respect to the ^{214}Ra isotope.

Given in Table VI are the isotope shifts between eight Ra isotopes for the $7s^2$ ^{1}S$_0$-$7s7p$ ^{3}P$_1$ intercombination (spin-forbidden) transition and the hyperfine structure constants A and B of the $7s7p$ ^{3}P$_1$ level for the odd isotopes. Here again, the isotope shifts are given with respect to the ^{214}Ra isotope.

In Table VII, isotope shifts of seven Ra isotopes for the $7s7p$ ^{3}P$_2$-$7s7d$ ^{3}D$_3$ transition at 644.6 nm and the hyperfine structure constants A and B of the $7s7p$ ^{3}P$_2$ and $7s7d$ ^{3}D$_3$ levels for the odd isotopes are given. The isotope shifts are given with respect to the ^{214}Ra isotope similar to the other transitions.

In Table VIII, the isotope shifts between ^{226}Ra and ^{225}Ra isotopes for the $7s6d$ ^{3}D$_1$-$7s7p$ ^{1}P$_1$ transition at 1428 nm and the hyperfine structure constants A and B of the $7s7p$ ^{1}P$_1$ and $7s6d$ ^{3}D$_1$ levels for the ^{225}Ra isotope are given.

III. RA II

The first experimental measurements on singly ionized Ra were conducted by Rasmussen in 1933 using an arc discharge as the light source. In total 62 lines were identified, corresponding to 43 energy levels. The energy levels are given in Table IX. In Table IX additional information on energy levels and their uncertainties is also given, which was obtained using the LOPT code[44]. In column 6 energy level values obtained with the LOPT code are given. Two uncertainties for the energy level values estimated by LOPT code are given; uncertainty 1 (Column 7) and uncertainty 2 (Column 8). Uncertainty 1 is the best determined separation from other energy levels and uncertainty 2 is determined for the excitation energy. Further, some of the energy levels given in Moore's compilation[9] are significantly improved by our level optimization, while a few were found to be much less accurate than implied by the number of significant figures given by Moore.

The ionization energy (IE) of Ra II has been redetermined because of the changes in the energy level values of Ra II. The series used to determine the IE is ng ^2G with $n = 5-11$, five members of which ($n = 5-9$) have been determined with uncertainties about 0.2 cm^{-1}. Using this series, two values of IE have been obtained; one by fitting a polarization formula and another by fitting a Ritz-type quantum defect formula as given in references[55,56]. The results of these two determinations are 81842.6(5) cm^{-1} and 81842.5(5) cm^{-1} resulting an average of 81842.5(5) cm^{-1}. One can also use the series ns $^2S_{1/2}$ ($n = 7-11$), nd $^2D_{3/2}$ ($n = 6-11$) and nd $^2D_{5/2}$ ($n = 6-12$) to determine IE. However, the IE limit values have larger scatter and uncertainties. Thus, the recommended IE value is 81842.5(5) cm^{-1}, which agrees with Rasmussen's value of 81842.31 cm^{-1} (as given by Moore) within the uncertainty.

In Table X intensity, observed wavelengths of the transitions, uncertainty of the observed wavelengths, observed wavenumbers, uncertainty of the observed wavenumbers, the corresponding lower and upper levels, line reference (Column 8), transition rates, uncertainty of the transition rates and the reference for the transition rates (Column 11) are given. Theoretical estimates of energy levels and dipole matrix elements for singly ionized Ra have been reported[32,51,57,58]. Here we present the transition rates from Sahoo *et al.*[32] as the uncertainty was given for the calculated values. In the intensity (Column 1), 'p' means 'perturbed by'. The corresponding perturbing species are given in parentheses. In the line reference (Column 8) '6' corresponds to the reference for the transition/line and the letters; HQ - Hilger Quartz, Q - Quartz prism, P - Glass prism and G - Grating (used in the red region) are the letter codes for the spectrometer used in the measurement. The Ritz wavelengths and their uncertainties for Ra II are given in Table XI. These values were calculated with the LOPT code[44]. For wavelengths less than 2000$\mathring{A}$ vacuum wavelengths are given and for wavelengths greater than 2000$\mathring{A}$ air wavelengths are given. The uncertainty of the given wavelengths mentioned in Ref. 6 is 0.1$\mathring{A}$ to 0.02$\mathring{A}$ between 3600 to 7000$\mathring{A}$ and 0.1$\mathring{A}$ between 1880 and 2200$\mathring{A}$[44]. In the red and near-infrared wavelength regions, i.e. 7000$\mathring{A}$ to 10000$\mathring{A}$ the uncertainty is 0.01$\mathring{A}$. Some of the observed wavenumbers given in column 4 differ from the wavenumbers given in the original work of Rasmussen[6]. The difference from the original values and the present reported values is due to the conversion formula used by Rasmussen for arriving at the wavenumbers from the measured wavelengths. Here we have used the three-term formula of Peck and Reeder[43] to arrive at the wavenumbers.

In Table XII, improved values of wavelengths in air, wavenumbers and uncertainty in wavenumbers are given for ^{212,213,214,225,226}Ra isotopes for the 7s $^2S_{1/2}$-

7s7p $^2P^0_{1/2,3/2}$ and 6d $^2D_{3/2}$-7s7p $^2P^0_{1/2,3/2}$ transitions. The wavenumbers are derived from the absolute frequency measurements of the corresponding transitions with a reference to a femtosecond optical frequency comb. The air wavelengths are calculated using the three-term formula (Equation 3) from Peck and Reeder[43].

Table XIII gives isotope shifts of 17 Ra isotopes for the 7s $^2S_{1/2}$ - 7p $^2P_{1/2}$ transition and of 8 isotopes for the 7s $^2S_{1/2}$ - 7p $^2P_{3/2}$ transition, 7 hyperfine structure constants for the 7s $^2S_{1/2}$ and 7p $^2P_{1/2}$ states, while for the 7p $^2P_{3/2}$ state, 3 magnetic dipole constants and 2 electric quadrupole constants measured by the ISOLDE group are given. The uncertainties are given in parentheses. The isotope shifts are given with respect to the ^{214}Ra isotope, which has a zero nuclear spin.

In Table XIV isotope shifts of six Ra isotopes for the 6d $^2D_{3/2}$ - 7p $^2P_{1/2}$ transition of singly ionized Ra and hyperfine structure constants of the 6d $^2D_{3/2}$ and 7p $^2P_{1/2}$ states measured at the University of Groningen are given.

Table XV consists of isotope shifts of the ^{212,213}Ra isotopes with respect to ^{214}Ra for the 6d $^2D_{3/2}$ - 7p $^2P_{3/2}$ transition and hyperfine structure of the 6d $^2D_{3/2}$ state. There is only one experimental measurement of lifetime of a state in singly ionized Ra. It is for the 6d $^2D_{5/2}$ state in ^{212}Ra with the lifetime of 232(4) ms (Table XVI). Although theoretical calculations are available for the lifetimes of different states in the literature[32,57–59], we have not included them in this compilation.

IV. SPECTROSCOPIC DATA

Ground state and configuration:
Ra I - $1s^2\ 2s^2 2p^6\ 3s^2 3p^6 3d^{10}\ 4s^2 4p^6 4d^{10} 4f^{14}\ 5s^2 5p^6 5d^{10}\ 6s^2 6p^6\ 7s^2\ {}^1S_0$

Ra II - $1s^2\ 2s^2 2p^6\ 3s^2 3p^6 3d^{10}\ 4s^2 4p^6 4d^{10} 4f^{14}\ 5s^2 5p^6 5d^{10}\ 6s^2 6p^6\ 7s\ {}^2S_{1/2}$

Ionization energies:
Ra I - 42573.36(2) cm^{-1}; 5.2784239(25) eV[17]
Ra II - 81842.5(5) cm^{-1}; 10.14718(6) eV

TABLE II: Energy levels of ^{226}Ra I.

Configuration	Term	J	Level (cm^{-1})	Uncertainty (cm^{-1})	Level LOPT (cm^{-1})	Uncertainty LOPT (cm^{-1})	Level reference
7s^2	^{1}S	0	0.0000		0.0000	0.0	
7s7p	^{3}P^0	0	13078.44		13078.40	0.04	9
	^{3}P^0	1	13999.357	0.001	13999.3569	0.0010	41
	^{3}P^0	2	16688.54		16688.54	0.04	9
7s6d	^{3}D	1	13715.85		13715.779	0.020	9
	^{3}D	2	13993.97		13993.94	0.04	9
	^{3}D	3	14707.35		14707.286	0.050	9
7s6d	^{1}D	2	17081.45		17081.313	0.060	9
7s7p	^{1}P^0	1	20715.6142	0.0005	20715.6142	0.0005	12
7s8s	^{3}S	1	26754.05		26754.015	0.030	9
7s8s	^{1}S	0					
6d7p	^{3}F^0	2	28038.05		28038.00	0.04	9
	^{3}F^0	3	30117.78		30117.72	0.06	9
	^{3}F^0	4	32367.78		32367.70	0.08	9
6d7p	^{1}D^0	2	30918.14		30918.01	0.07	9
7s8p	^{3}P^0	0	31085.88		31085.80	0.15	9
	^{3}P^0	1	31563.29		31563.40	0.13	9
	^{3}P^0	2	31874.44		31874.25	0.11	9
7p^2 (6d^2)^{51}	^{3}P	0					9
	^{3}P	1	31248.61		31248.61	0.07	9
	^{3}P	2	32941.13		32941.20	0.03	9
7s7d	^{3}D	1	32000.82		32000.78	0.05	9
	^{3}D	2	31993.41		31993.40	0.06	9
	^{3}D	3	32197.28		32197.27	0.07	9
7p^2 (7s7d)^{21}, (6d^2)^{51}	^{1}D	2	32214.84		32214.76	0.06	9
6d7p	^{3}D^0	1	32229.97		32229.89	0.06	9
	^{3}D^0	2	32506.59		32506.52	0.05	9
	^{3}D^0	3	33197.46		33197.39	0.07	9
7s8p	^{1}P^0	1	32857.68		32857.54	0.08	9
6d7p	^{3}P^0	0	33782.41		33782.32	0.20	9
	^{3}P^0	1	33823.70		33823.63	0.15	9
	^{3}P^0	2	34382.91		34382.83	0.08	9

TABLE II Energy levels, ^{226}Ra I – continued

Configuration	Term	J	Level (cm^{-1})	Uncertainty (cm^{-1})	Level LOPT (cm^{-1})	Uncertainty LOPT (cm^{-1})	Level reference
7s9s	^{3}S	1	34475.78		34475.74	0.12	9
7s5f	^{3}F^0	2	35255.65		35255.51	0.09	9
	^{3}F^0	3	35268.02		35267.88	0.09	9
	^{3}F^0	4	35294.40		35294.32	0.10	9
7s6f	^{3}F^0	2					9
	^{3}F^0	3	37922.26		37922.2	0.3	9
	^{3}F^0	4	37929.64		37929.6	0.3	9
7s7f	^{3}F^0	2					9
	^{3}F^0	3	39360.96		39360.9	0.3	9
	^{3}F^0	4	39366.98		39366.9	0.3	9
7s8f	^{3}F^0	2	40219.5		40219.4	0.4	9
	^{3}F^0	3	40219.5		40219.5	0.4	9
	^{3}F^0	4	40231.58		40231.5	0.3	9
7s13p	^{1}P^0	1	41034.401	0.006	41034.400	0.017	17
7s14p	^{1}P^0	1	41338.940	0.006	41338.940	0.017	17
7s15p	^{1}P^0	1	41562.296	0.006	41562.296	0.017	17
7s16p	^{1}P^0	1	41730.581	0.006	41730.580	0.017	17
7s17p	^{1}P^0	1	41860.241	0.006	41860.240	0.018	17
7s18p	^{1}P^0	1	41962.282	0.006	41962.281	0.018	17
7s19p	^{1}P^0	1	42043.869	0.006	42043.868	0.018	17
7s20p	^{1}P^0	1	42110.347	0.006	42110.347	0.018	17
7s21p	^{1}P^0	1	42164.971	0.006	42164.971	0.018	17
7s22p	^{1}P^0	1	42210.520	0.006	42210.520	0.018	17
7s23p	^{1}P^0	1	42248.829	0.006	42248.829	0.018	17
7s24p	^{1}P^0	1	42281.384	0.006	42281.383	0.018	17
7s25p	^{1}P^0	1	42309.321	0.006	42309.320	0.018	17
7s26p	^{1}P^0	1	42333.336	0.006	42333.335	0.018	17
7s27p	^{1}P^0	1	42354.358	0.006	42354.357	0.018	17
7s28p	^{1}P^0	1	42372.773	0.006	42372.773	0.018	17
7s29p	^{1}P^0	1	42388.773	0.006	42388.773	0.018	17
7s30p	^{1}P^0	1	42403.096	0.006	42403.096	0.018	17
7s31p	^{1}P^0	1	42415.790	0.006	42415.789	0.018	17
7s32p	^{1}P^0	1	42427.050	0.006	42427.050	0.018	17
7s33p	^{1}P^0	1	42437.097	0.006	42437.097	0.018	17
7s34p	^{1}P^0	1	42446.203	0.006	42446.203	0.018	17
7s35p	^{1}P^0	1	42454.517	0.006	42454.516	0.018	17
7s36p	^{1}P^0	1	42461.863	0.006	42461.863	0.018	17
7s37p	^{1}P^0	1	42468.788	0.006	42468.788	0.018	17
7s38p	^{1}P^0	1	42474.918	0.006	42474.917	0.018	17
7s39p	^{1}P^0	1	42480.501	0.006	42480.501	0.018	17
7s40p	^{1}P^0	1	42485.662	0.006	42485.662	0.018	17
7s41p	^{1}P^0	1	42490.375	0.006	42490.375	0.018	17
7s42p	^{1}P^0	1	42494.765	0.006	42494.765	0.018	17
7s43p	^{1}P^0	1	42498.882	0.006	42498.882	0.018	17
7s44p	^{1}P^0	1	42502.575	0.006	42502.575	0.018	17
7s45p	^{1}P^0	1	42506.119	0.006	42506.118	0.018	17
7s46p	^{1}P^0	1	42509.289	0.006	42509.288	0.018	17
7s47p	^{1}P^0	1	42512.260	0.006	42512.260	0.018	17
7s48p	^{1}P^0	1	42515.131	0.006	42515.131	0.018	17
7s49p	^{1}P^0	1	42517.803	0.006	42517.803	0.018	17
7s50p	^{1}P^0	1	42520.126	0.006	42520.125	0.018	17
7s51p	^{1}P^0	1	42522.424	0.006	42522.423	0.018	17
7s52p	^{1}P^0	1	42524.473	0.006	42524.473	0.018	17
Ra II ($6p^6$ 7s $^2S_{1/2}$)	Limit		42573.36	0.02			17

TABLE III: Wavelengths, Wave numbers, classification and transition probabilities in neutral radium (^{226}Ra), Ra I

Intensity	Observed $\lambda_{air}/\AA$	Unc. Obs. $\lambda_{air}/\AA$	Obs. σ/cm^{-1}	Lower level	Upper level	Line reference(s)	Ritz $\lambda/\AA$	Unc. Ritz $\lambda/\AA$	A_{ik}/s^{-1}	Reference
	2350.8675	0.0010	42524.473	$7s^2\ ^1S_0$	$7s52p\ ^1P_1^0$	17				
	2350.9808	0.0010	42522.423	$7s^2\ ^1S_0$	$7s51p\ ^1P_1^0$	17				
	2351.1079	0.0010	42520.125	$7s^2\ ^1S_0$	$7s50p\ ^1P_1^0$	17				
	2351.2363	0.0010	42517.803	$7s^2\ ^1S_0$	$7s49p\ ^1P_1^0$	17				
	2351.3841	0.0010	42515.131	$7s^2\ ^1S_0$	$7s48p\ ^1P_1^0$	17				
	2351.5429	0.0010	42512.260	$7s^2\ ^1S_0$	$7s47p\ ^1P_1^0$	17				
	2351.7073	0.0010	42509.288	$7s^2\ ^1S_0$	$7s46p\ ^1P_1^0$	17				
	2351.8827	0.0010	42506.118	$7s^2\ ^1S_0$	$7s45p\ ^1P_1^0$	17				
	2352.0788	0.0010	42502.575	$7s^2\ ^1S_0$	$7s44p\ ^1P_1^0$	17				
	2352.2832	0.0010	42498.882	$7s^2\ ^1S_0$	$7s43p\ ^1P_1^0$	17				
	2352.5111	0.0010	42494.765	$7s^2\ ^1S_0$	$7s42p\ ^1P_1^0$	17				
	2352.7542	0.0010	42490.375	$7s^2\ ^1S_0$	$7s41p\ ^1P_1^0$	17				
	2353.0152	0.0010	42485.662	$7s^2\ ^1S_0$	$7s40p\ ^1P_1^0$	17				
	2353.3011	0.0010	42480.501	$7s^2\ ^1S_0$	$7s39p\ ^1P_1^0$	17				
	2353.6105	0.0010	42474.917	$7s^2\ ^1S_0$	$7s38p\ ^1P_1^0$	17				
	2353.9502	0.0010	42468.788	$7s^2\ ^1S_0$	$7s37p\ ^1P_1^0$	17				
	2354.3341	0.0010	42461.863	$7s^2\ ^1S_0$	$7s36p\ ^1P_1^0$	17				
	2354.7416	0.0010	42454.516	$7s^2\ ^1S_0$	$7s35p\ ^1P_1^0$	17				
	2355.2028	0.0010	42446.203	$7s^2\ ^1S_0$	$7s34p\ ^1P_1^0$	17				
	2355.7082	0.0010	42437.097	$7s^2\ ^1S_0$	$7s33p\ ^1P_1^0$	17				
	2356.2661	0.0010	42427.050	$7s^2\ ^1S_0$	$7s32p\ ^1P_1^0$	17				
	2356.8917	0.0010	42415.789	$7s^2\ ^1S_0$	$7s31p\ ^1P_1^0$	17				
	2357.5973	0.0010	42403.096	$7s^2\ ^1S_0$	$7s30p\ ^1P_1^0$	17				
	2358.3940	0.0010	42388.773	$7s^2\ ^1S_0$	$7s29p\ ^1P_1^0$	17				
	2359.2846	0.0010	42372.773	$7s^2\ ^1S_0$	$7s28p\ ^1P_1^0$	17				
	2360.3105	0.0010	42354.357	$7s^2\ ^1S_0$	$7s27p\ ^1P_1^0$	17				
	2361.4827	0.0010	42333.335	$7s^2\ ^1S_0$	$7s26p\ ^1P_1^0$	17				
	2362.8232	0.0010	42309.320	$7s^2\ ^1S_0$	$7s25p\ ^1P_1^0$	17				
	2364.3845	0.0010	42281.383	$7s^2\ ^1S_0$	$7s24p\ ^1P_1^0$	17				
	2366.2065	0.0010	42248.829	$7s^2\ ^1S_0$	$7s23p\ ^1P_1^0$	17				
	2368.3542	0.0010	42210.520	$7s^2\ ^1S_0$	$7s22p\ ^1P_1^0$	17				
	2370.9128	0.0010	42164.971	$7s^2\ ^1S_0$	$7s21p\ ^1P_1^0$	17				
	2373.9885	0.0010	42110.347	$7s^2\ ^1S_0$	$7s20p\ ^1P_1^0$	17				
	2377.7425	0.0010	42043.868	$7s^2\ ^1S_0$	$7s19p\ ^1P_1^0$	17				
	2382.3659	0.0010	41962.281	$7s^2\ ^1S_0$	$7s18p\ ^1P_1^0$	17				
	2388.1738	0.0010	41860.240	$7s^2\ ^1S_0$	$7s17p\ ^1P_1^0$	17				
	2395.5946	0.0010	41730.580	$7s^2\ ^1S_0$	$7s16p\ ^1P_1^0$	17				
	2405.2950	0.0010	41562.296	$7s^2\ ^1S_0$	$7s15p\ ^1P_1^0$	17				
	2418.2919	0.0010	41338.940	$7s^2\ ^1S_0$	$7s14p\ ^1P_1^0$	17				
	2436.2408	0.0010	41034.400	$7s^2\ ^1S_0$	$7s13p\ ^1P_1^0$	17				
2	2955.65	0.05	33823.6	$7s^2\ ^1S_0$	$6d7p\ ^3P_1^0$	8	2955.649	0.012		
			32857.68	$7s^2\ ^1S_0$	$7s8p\ ^1P_1^0$	9	3042.557	0.008		
5	3101.80	0.05	32230.0	$7s^2\ ^1S_0$	$6d7p\ ^3D_1^0$	7, 8	3101.810	0.005		
3	3771.57	0.05	26506.66			7				
1	3772.00	0.05	26503.6	$7s6d\ ^3D_1$	$7s8f\ ^3F_2^0$	7				
1	3812.00	0.05	26225.5	$7s6d\ ^3D_2$	$7s8f\ ^3F_3^0$	7				
1	3916.74	0.05	25524.2	$7s6d\ ^3D_3$	$7s8f\ ^3F_4^0$	7				
1	3941.02	0.05	25367.0	$7s6d\ ^3D_2$	$7s7f\ ^3F_3^0$	7				
1	4010.30	0.05	24928.77			7				
1	4054.07	0.05	24659.6	$7s6d\ ^3D_3$	$7s7f\ ^3F_4^0$	7				
4	4177.98	0.05	23928.3	$7s6d\ ^3D_2$	$7s6f\ ^3F_3^0$	7				
3	4265.12	0.05	23439.43			7				
4	4305.00	0.05	23222.3	$7s6d\ ^3D_3$	$7s6f\ ^3F_4^0$	7				
2	4366.30	0.05	22896.27			7				
2	4426.35	0.05	22585.65			7				
1	4444.47	0.05	22493.57			7				
8	4641.290	0.020	21539.70	$7s6d\ ^3D_1$	$7s5f\ ^3F_2^0$	7	4641.284	0.019		

TABLE III Wavelengths, Wave numbers, classification and transition probabilities in ^{226}Ra I – continued

Intensity	Observed $\lambda_{air}/\AA$	Unc. Obs. $\lambda_{air}/\AA$	Observed σ/cm^{-1}	Lower level	Upper level	Line reference(s)	Ritz $\lambda/\AA$	Unc. Ritz $\lambda/\AA$	A_{ik}/s^{-1}	Reference
1	4672.19	0.05	21397.25	$7s7p\ ^3\mathrm{P}^0_0$	$7s9s\ ^3\mathrm{S}_1$	7	4672.17	0.03		
8	4699.280	0.020	21273.90	$7s6d\ ^3\mathrm{D}_2$	$7s5f\ ^3\mathrm{F}^0_3$	7	4699.272	0.018		
2	4701.97	0.05	21261.73	$7s6d\ ^3\mathrm{D}_2$	$7s5f\ ^3\mathrm{F}^0_2$	7	4702.006	0.020		
2	4740.07	0.05	21090.86			7				
3	4803.11	0.05	20814.04			7				
100	4825.92806	0.00012	20715.6142	$7s^2\ ^1\mathrm{S}_0$	$7s7p\ ^1\mathrm{P}^0_1$	7, 12	4825.92805	0.0012	1.80×10^8	53
3	4837.27	0.05	20667.04	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{P}^0_2$	7	4837.268	0.017		
10	4856.070	0.020	20587.03	$7s6d\ ^3\mathrm{D}_3$	$7s5f\ ^3\mathrm{F}^0_4$	7				
2	4862.27	0.05	20560.78	$7s6d\ ^3\mathrm{D}_3$	$7s5f\ ^3\mathrm{F}^0_3$	7	4862.314	0.021		
3	4882.28	0.05	20476.52	$7s7p\ ^3\mathrm{P}^0_1$	$7s9s\ ^3\mathrm{S}_1$	7	4882.31	0.03		
4	4903.24	0.05	20388.98	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{P}^0_2$	7	4903.263	0.017		
5	4971.77	0.05	20107.95	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{P}^0_1$	7, 8	4971.79	0.03		
4	4982.03	0.05	20066.54	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{P}^0_0$	7, 8				
5	5041.56	0.05	19829.60	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{P}^0_1$	7, 8	5041.54	0.04		
6	5081.03	0.05	19675.56	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^3\mathrm{P}^0_2$	7	5081.035	0.017		
10	5097.56	0.05	19611.78			7				
10	5205.930	0.020	19203.52	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{D}^0_3$	7	5205.948	0.015		
4	5263.96	0.05	18991.83			7				
2	5277.91	0.05	18941.62	$7s7p\ ^3\mathrm{P}^0_1$	$7p^2\ ^3\mathrm{P}_2$	7	5277.849	0.008		
10	5283.280	0.020	18922.37	$7s7p\ ^3\mathrm{P}^0_0$	$7s7d\ ^3\mathrm{D}_1$	7	5283.277	0.014		
10	5320.290	0.020	18790.74	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{D}^0_2$	7	5320.290	0.014		
10	5399.800	0.020	18514.06	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{D}^0_1$	7	5399.784	0.014		
20	5400.230	0.020	18512.58	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{D}^0_2$	7	5400.231	0.013		
20	5406.810	0.020	18490.05	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^3\mathrm{D}^0_3$	7	5406.795	0.016		
8	5482.130	0.020	18236.02	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{D}^0_1$	7	5482.150	0.015		
4	5488.32	0.05	18215.45	$7s7p\ ^3\mathrm{P}^0_1$	$7p^2\ ^1\mathrm{D}_2$	7	5488.334	0.017		
2	5497.83	0.05	18183.95			7				
10	5501.980	0.020	18170.23	$7s7p\ ^3\mathrm{P}^0_0$	$7p^2\ ^3\mathrm{P}_1$	7	5501.985	0.018		
4	5505.50	0.05	18158.61	$7s6d\ ^3\mathrm{D}_1$	$7s8p\ ^3\mathrm{P}^0_2$	7, 8	5505.542	0.024		
3	5544.25	0.05	18031.71			7				
10	5553.570	0.020	18001.44	$7s7p\ ^3\mathrm{P}^0_1$	$7s7d\ ^3\mathrm{D}_1$	7	5553.574	0.015		
20	5555.850	0.020	17994.05	$7s7p\ ^3\mathrm{P}^0_1$	$7s7d\ ^3\mathrm{D}_2$	7	5555.852	0.018		
2	5591.15	0.05	17880.44	$7s6d\ ^3\mathrm{D}_2$	$7s8p\ ^3\mathrm{P}^0_2$	7, 8	5591.192	0.024		
5	5601.50	0.05	17847.41	$7s6d\ ^3\mathrm{D}_1$	$7s8p\ ^3\mathrm{P}^0_1$	7	5601.43	0.03		
10	5616.660	0.020	17799.23	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^3\mathrm{D}^0_2$	7	5616.660	0.015		
3	5620.47	0.05	17787.17	$7s7p\ ^3\mathrm{P}^0_2$	$7s9s\ ^3\mathrm{S}_1$	7	5620.46	0.04		
50	5660.810	0.020	17660.42	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^3\mathrm{F}^0_4$	7			4.30×10^7	53
3	5690.16	0.05	17569.32	$7s6d\ ^3\mathrm{D}_2$	$7s8p\ ^3\mathrm{P}^0_1$	7	5690.12	0.03		
4	5755.45	0.05	17370.02	$7s6d\ ^3\mathrm{D}_1$	$7s8p\ ^3\mathrm{P}^0_0$	7, 8				
5	5778.28	0.05	17301.39	$7s6d\ ^1\mathrm{D}_2$	$6d7p\ ^3\mathrm{P}^0_2$	7	5778.24	0.03		
5	5795.78	0.05	17249.16	$7s7p\ ^3\mathrm{P}^0_1$	$7p^2\ ^3\mathrm{P}_1$	7	5795.745	0.023		
5	5811.58	0.05	17202.26	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^1\mathrm{D}^0_2$	7	5811.588	0.021	4.00×10^6	53
3	5823.49	0.05	17167.07	$7s6d\ ^3\mathrm{D}_3$	$7s8p\ ^3\mathrm{P}^0_2$	7, 8	5823.53	0.03		
2	5907.20	0.05	16923.80	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^1\mathrm{D}^0_2$	7	5907.107	0.021	5.50×10^4	53
2	5957.17	0.05	16781.86			7				
5	5957.67	0.05	16780.45			7				
3	6151.19	0.05	16252.52	$7s7p\ ^3\mathrm{P}^0_2$	$7p^2\ ^3\mathrm{P}_2$	7	6151.137	0.019		
5	6167.03	0.05	16210.77	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^1\mathrm{D}^0_2$	7	6167.049	0.024	1.50×10^7	53
30	6200.300	0.020	16123.79	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{F}^0_3$	7	6200.304	0.016	2.50×10^7	53
10	6336.900	0.020	15776.22	$7s6d\ ^1\mathrm{D}_2$	$7s8p\ ^1\mathrm{P}^0_1$	7				
3	6438.90	0.05	15526.31	$7s7p\ ^3\mathrm{P}^0_2$	$7p^2\ ^1\mathrm{D}_2$	7	6438.94	0.03		
20	6446.200	0.020	15508.73	$7s7p\ ^3\mathrm{P}^0_2$	$7s7d\ ^3\mathrm{D}_3$	7				
2	6474.47	0.05	15441.02			7				
20	6487.320	0.020	15410.43	$7s6d\ ^3\mathrm{D}_3$	$6d7p\ ^3\mathrm{F}^0_3$	7	6487.317	0.017	9.90×10^6	53
2	6528.92	0.05	15312.24	$7s7p\ ^3\mathrm{P}^0_2$	$7s7d\ ^3\mathrm{D}_1$	7	6528.92	0.02		
3	6532.08	0.05	15304.83	$7s7p\ ^3\mathrm{P}^0_2$	$7s7d\ ^3\mathrm{D}_2$	7	6532.07	0.03		
3	6545.93	0.05	15272.46			7				
4	6585.41	0.05	15180.90			7				
3	6599.47	0.05	15148.55	$7s6d\ ^1\mathrm{D}_2$	$6d7p\ ^3\mathrm{D}^0_1$	7	6599.46	0.03		

TABLE III Wavelengths, Wave numbers, classification and transition probabilities in ^{226}Ra I – continued

Intensity	Observed $\lambda_{air}/\mathring{A}$	Unc. Obs. $\lambda_{air}/\mathring{A}$	Observed σ/cm^{-1}	Lower level	Upper level	Line reference(s)	Ritz $\lambda/\mathring{A}$	Unc. Ritz $\lambda/\mathring{A}$	A_{ik}/s^{-1}	Reference
2	6645.95	0.05	15042.61			7				
2	6653.33	0.05	15025.92			7				
4	6758.20	0.05	14792.76	$7s6d\ ^1\mathrm{D}_2$	$7s8p\ ^3\mathrm{P}^0_2$	7, 8	6758.12	0.04		
3	6903.10	0.05	14482.25	$7s6d\ ^1\mathrm{D}_2$	$7s8p\ ^3\mathrm{P}^0_1$	7	6903.18	0.04		
20	6980.220	0.020	14322.25	$7s6d\ ^3\mathrm{D}_1$	$6d7p\ ^3\mathrm{F}^0_2$	7	6980.232	0.017	1.70×10^7	53
20	7118.500	0.020	14044.03	$7s6d\ ^3\mathrm{D}_2$	$6d7p\ ^3\mathrm{F}^0_2$	7	7118.486	0.017	9.80×10^6	53
50	7141.2167	0.0005	13999.3569	$7s^2\ ^1\mathrm{S}_0$	$7s7p\ ^3\mathrm{P}^0_1$	7, 41	7141.2167	0.0005	2.37×10^6	41[60]
20	7225.160	0.020	13836.71	$7s6d\ ^1\mathrm{D}_2$	$6d7p\ ^1\mathrm{D}^0_2$	7	7225.167	0.019	3.50×10^7	53
10	7310.270	0.020	13675.62	$7s7p\ ^3\mathrm{P}^0_0$	$7s8s\ ^3\mathrm{S}_1$	7	7310.271	0.018	8.50×10^6	53
1	7468.21	0.05	13386.41			7				
2	7499.87	0.05	13329.90			7				
2	7565.49	0.05	13214.31			7				
1	7836.09	0.05	12757.96			7				
20	7838.120	0.020	12754.65	$7s7p\ ^3\mathrm{P}^0_1$	$7s8s\ ^3\mathrm{S}_1$	7	7838.116	0.018	2.10×10^7	53
2	7877.08	0.05	12691.57			7				
3	7896.43	0.05	12660.47			7				
3	8005.13	0.05	12488.56			7				
2	8019.25	0.05	12466.58			7				
6	8177.310	0.020	12225.60	$7s7p\ ^1\mathrm{P}^0_1$	$7p^2\ ^3\mathrm{P}_2$	7	8177.319	0.019		
3	8248.70	0.05	12119.79			7				
3	8269.03	0.05	12090.00			7				
5	8335.07	0.05	11994.21			7				
4	8693.94	0.05	11499.11	$7s7p\ ^1\mathrm{P}^0_1$	$7p^2\ ^1\mathrm{D}_2$	7	8693.91	0.04		
2	9094.80	0.05	10992.28			7				
5	9932.21	0.05	10065.49	$7s7p\ ^3\mathrm{P}^0_2$	$7s8s\ ^3\mathrm{S}_1$	7	9932.23	0.04	2.50×10^7	53
	14282.14	0.04	6999.84	$7s6d\ ^3\mathrm{D}_1$	$7s7p\ ^1\mathrm{P}^0_1$	10	14286.05	0.04	4.40×10^4	53

TABLE IV. Lifetimes of energy levels - ^{226}Ra I

Level	7s7p $^3P_1^0$	7s6d $^1D_2$	7s6d $^3D_1$
Lifetime	422(20) ns	385(45) μs	510(60) μs
Reference	41	42	10

TABLE V. Isotope shifts and hyperfine structure constants A and B of 7s^2 $^1S_0$ and 7s7p $^1P_1^0$ levels in Ra I

	7s^2 $^1S_0$	7s7p $^1P_1^0$			
Atomic mass		Isotopic frequency shift (MHz)	HFS A (MHz)	HFS B (MHz)	Reference
208		8485(8)			21
209		8313(8)	-656.5(1.2)	135.0(5.0)	21
210		6003(6)			21
211		5522(4)	-668.4(0.9)	162.0(4.0)	21
212		3256(3)			21
213		2167(3)	-2314.8(2.5)		21
214	0	0			21
220		-21904(8)			21
221		-25945(8)	136.3(0.8)	669.6(3.0)	21
222		-28827(8)			21
223		-32453(9)	-344.5(0.9)	421.5(1.6)	21
224		-35123(9)			21
225		-38888(10)	2796.5(2.5)		21
226		-41124(11)			21
227		-43937(12)	512.5(2.3)	531.0(5.0)	21
228		-46950(12)			21
229		-50066(15)	-382.7(1.4)	1042.0(5.0)	21
230		-53634(15)			21
232		-59585(17)			21

TABLE VI. Isotope shifts and hyperfine structure constants A and B of 7s^2 $^1S_0$ and 7s7p $^3P_1^0$ levels in Ra I

	7s^2 $^1S_0$	7s7p $^3P_1^0$			
Atomic mass		Isotopic frequency shift (MHz)	HFS A (MHz)	HFS B (MHz)	Reference
212		3303(5)			21
214	0	0			21
221			-475.8(1.4)	-747.6(2.6)	21
222		-29253(9)			21
223		-32934(10)	1,202.1(0.6)	-470.2(1.2)	21
224		-35644(11)			21
225		-39465(12)	-9,793.9(4.3)		21
226		-41733(12)			21

TABLE VII. Isotope shifts and hyperfine structure constants A and B of the 7s7p $^3P_2^0$ level and the 7s7d $^3D_3$ level in Ra I

	7s7p $^3P_2^0$		7s7d $^3D_3$			
Atomic mass	HFS A(MHz)	HFS B (MHz)	Isotopic frequency shift (MHz)	HFS A (MHz)	HFS B (MHz)	Reference
212			-887(4)			21
214			0			21
221	278.6(0.9)	1103.8(3.9)	7001(7)	213.3(0.6)	117.8(4.4)	21
222			7781(4)			21
223	699.6(3.3)	688.5(4.0)	8762(6)	535.6(4.6)	73.9(3.0)	21
224			9481(5)			21
226			11102(6)			21

TABLE VIII. Isotope shifts and hyperfine structure constants A and B of the 7s7p $^1P_1^0$ level and the 7s6d $^3D_1$ level in Ra I

	7s7p $^1P_1^0$		7s6d $^3D_1$			
Atomic mass	HFS A(MHz)	HFS B (MHz)	Isotopic frequency shift (MHz)	HFS A (MHz)	HFS B (MHz)	Reference
225	2797.3 (1.5)		540.2 (2.0)	4687.7 (1.5)		10
226			0			10

TABLE X: Wavelengths, Wave numbers, classification and transition probabilities for observed lines in radium ion, ^{226}Ra II

Intensity[61]	Obs. $\lambda/\AA$	Unc. Obs. $\lambda/\AA$	Obs. σ/cm^{-1}	Unc. Obs. σ/cm^{-1}	Lower level	Upper level	Line ref.[62]	A_{ik}/s^{-1}	Acc.[63]	Ref.
5	1888.70	0.20	52946		6d ^{2}D$_{5/2}$	7f ^{2}F$^0_{7/2}$	6HQ			
8	1908.70	0.20	52392		7s ^{2}S$_{1/2}$	8p ^{2}P$^0_{3/2}$	6HQ	1.75×10^7	B	32
4	1972.60	0.20	50695		7p ^{2}P$^0_{1/2}$	10d ^{2}D$_{3/2}$	6HQ			
2	1976.00	0.20	50607		7s ^{2}S$_{1/2}$	8p ^{2}P$^0_{1/2}$	6HQ			
2	2006.40	0.20	49824		7p ^{2}P$^0_{1/2}$	11s ^{2}S$_{1/2}$	6HQ			
3	2012.75	0.10	49667		6d ^{2}D$_{5/2}$	9p ^{2}P$^0_{3/2}$	6HQ			
2	2070.60	0.20	48280		7p ^{2}P$^0_{3/2}$	11d ^{2}D$_{5/2}$	6HQ			
4	2107.60	0.20	47432		6d ^{2}D$_{3/2}$	6f ^{2}F$^0_{5/2}$	6Q	2.67×10^7	B	32
5	2131.00	0.20	46912		7p ^{2}P$^0_{1/2}$	9d ^{2}D$_{3/2}$	6Q			
10	2169.90	0.20	46071		6d ^{2}D$_{5/2}$	6f ^{2}F$^0_{7/2}$	6Q			
4	2177.30	0.20	45914		7p ^{2}P$^0_{3/2}$	10d ^{2}D$_{5/2}$	6Q			
1	2181.10	0.20	45834		7p ^{2}P$^0_{3/2}$	10d ^{2}D$_{3/2}$	6Q			
2	2197.80	0.20	45486		7p ^{2}P$^0_{1/2}$	10s ^{2}S$_{1/2}$	6Q			
3	2223.30	0.20	44964		7p ^{2}P$^0_{3/2}$	11s ^{2}S$_{1/2}$	6Q			
8	2369.730	0.020	42186.0		7p ^{2}P$^0_{3/2}$	9d ^{2}D$_{5/2}$	6Q			
3	2377.100	0.020	42055.2		7p ^{2}P$^0_{3/2}$	9d ^{2}D$_{3/2}$	6Q			
8	2460.55	0.03	40629.0		7p ^{2}P$^0_{3/2}$	10s ^{2}S$_{1/2}$	6Q			
10	2475.50	0.03	40383.7		7p ^{2}P$^0_{1/2}$	8d ^{2}D$_{3/2}$	6Q			
4	2480.11	0.03	40308.6		6d ^{2}D$_{3/2}$	8p ^{2}P$^0_{3/2}$	6Q	7.5×10^5	C	32
8	2586.61	0.03	38649.1		6d ^{2}D$_{5/2}$	8p ^{2}P$^0_{3/2}$	6Q			
2	2595.15	0.03	38521.9		6d ^{2}D$_{3/2}$	8p ^{2}P$^0_{1/2}$	6Q			
10	2643.73	0.03	37814.1		7p ^{2}P$^0_{1/2}$	9s ^{2}S$_{1/2}$	6Q			
20	2708.960	0.020	36903.6		6d ^{2}D$_{3/2}$	5f ^{2}F$^0_{5/2}$	6G	3.70×10^8	A	32
10p (Mg II)	2795.21	0.07	35764.9		7p ^{2}P$^0_{3/2}$	8d ^{2}D$_{5/2}$	6Q			
30	2813.760	0.020	35529.2		6d ^{2}D$_{5/2}$	5f ^{2}F$^0_{7/2}$	6G			
6	2836.460	0.020	35244.85		6d ^{2}D$_{5/2}$	5f ^{2}F$^0_{5/2}$	6G			
10	3033.44	0.04	32956.3		7p ^{2}P$^0_{3/2}$	9s ^{2}S$_{1/2}$	6Q			
2	3423.10	0.20	29204.9		5f ^{2}F$^0_{5/2}$	11g ^{2}G$_{7/2,9/2}$	6Q			
2	3456.20	0.20	28925.2		5f ^{2}F$^0_{7/2}$	11g ^{2}G$_{7/2,9/2}$	6Q			
2	3514.80	0.20	28443.0		5f ^{2}F$^0_{5/2}$	10g ^{2}G$_{7/2,9/2}$	6Q			
2	3550.20	0.20	28159.4		5f ^{2}F$^0_{7/2}$	10g ^{2}G$_{7/2,9/2}$	6Q			
100	3649.550	0.020	27392.83		7p ^{2}P$^0_{1/2}$	7d ^{2}D$_{3/2}$	6G			
3	3686.160	0.020	27120.78		5f ^{2}F$^0_{7/2}$	9g ^{2}G$_{7/2,9/2}$	6P			
200	3814.4219	0.0003	26208.8538	0.0021	7s ^{2}S$_{1/2}$	7p ^{2}P$^0_{3/2}$	6G, 13	1.880×10^8	A	32
5p (Cl II)	3851.90	0.05	25953.9		5f ^{2}F$^0_{5/2}$	8g ^{2}G$_{7/2,9/2}$	6P			
5	3894.55	0.03	25669.63		5f ^{2}F$^0_{7/2}$	8g ^{2}G$_{7/2,9/2}$	6P			
8	4194.09	0.03	23836.36		5f ^{2}F$^0_{5/2}$	7g ^{2}G$_{7/2,9/2}$	6P			
2	4195.56	0.03	23828.01		8p ^{2}P$^0_{1/2}$	11d ^{2}D$_{3/2}$	6P			
1	4220.24	0.03	23688.66		8p ^{2}P$^0_{3/2}$	12d ^{2}D$_{5/2}$	6P			
8	4244.72	0.04	23552.05		5f ^{2}F$^0_{7/2}$	7g ^{2}G$_{7/2,9/2}$	6P			
100p (H I)	4340.64	0.07	23031.6		7p ^{2}P$^0_{3/2}$	7d ^{2}D$_{5/2}$	6P			
20	4436.270	0.020	22535.13		7p ^{2}P$^0_{3/2}$	7d ^{2}D$_{3/2}$	6G			
30	4533.110	0.020	22053.73		7p ^{2}P$^0_{1/2}$	8s ^{2}S$_{1/2}$	6G			
3	4663.52	0.05	21437.03		8p ^{2}P$^0_{1/2}$	10d ^{2}D$_{3/2}$	6P			
100	4682.23940	0.00017	21351.3259	0.0008	7s ^{2}S$_{1/2}$	7p ^{2}P$^0_{1/2}$	6G, 13	1.061×10^8	A	32
10	4859.41	0.05	20572.88		5f ^{2}F$^0_{5/2}$	6g ^{2}G$_{7/2,9/2}$	6P			
10	4927.53	0.05	20288.48		5f ^{2}F$^0_{7/2}$	6g ^{2}G$_{7/2,9/2}$	6P			
3	4997.26	0.05	20005.39		8s ^{2}S$_{1/2}$	9p ^{2}P$^0_{3/2}$	6P			
3	5066.57	0.05	19731.72		8p ^{2}P$^0_{3/2}$	10d ^{2}D$_{5/2}$	6P			
1	5323.09	0.06	18780.86		8p ^{2}P$^0_{3/2}$	11s ^{2}S$_{1/2}$	6P			
3	5623.43	0.07	17777.81		7d ^{2}D$_{3/2}$	7f ^{2}F$^0_{5/2}$	6P			
6	5661.73	0.07	17657.55		8p ^{2}P$^0_{1/2}$	9d ^{2}D$_{3/2}$	6P			
4	5728.83	0.07	17450.73		7d ^{2}D$_{5/2}$	7f ^{2}F$^0_{7/2}$	6P			

TABLE X Wavelengths, Wave numbers, classification and transition probabilities in radium ion, ^{226}Ra II– continued

Intensity	Obs. $\lambda_{air}/\AA$	Unc. Obs. $\lambda_{air}/\AA$	Obs. σ/cm^{-1}	Unc. Obs. σ/cm^{-1}	Lower level	Upper level	Line ref.	A_{ik}/s^{-1}	Acc.	Ref.
20	5813.630	0.020	17196.19		7p $^2P^0_{3/2}$	8s $^2S_{1/2}$	6G			
1	6158.73	0.08	16232.62		8p $^2P^0_{1/2}$	10s $^2S_{1/2}$	6P			
4	6247.16	0.08	16002.85		8p $^2P^0_{3/2}$	9d $^2D_{5/2}$	6P			
1	6298.56	0.08	15872.26		8p $^2P^0_{3/2}$	9d $^2D_{3/2}$	6P			
10	6593.340	0.020	15162.63		5f $^2F^0_{5/2}$	5g $^2G_{7/2,9/2}$	6G			
10	6719.320	0.020	14878.35		5f $^2F^0_{7/2}$	5g $^2G_{7/2,9/2}$	6G			
1	6920.11	0.10	14446.65		8p $^2P^0_{3/2}$	10s $^2S_{1/2}$	6P			
5	7078.02	0.10	14124.35		6d $^2D_{3/2}$	7p $^2P^0_{3/2}$	6G	3.39×10^6	B	32
50	8019.700	0.020	12465.87		6d $^2D_{5/2}$	7p $^2P^0_{3/2}$	6G	2.29×10^7	B	32
	10787.9609	0.0007	9267.0538	0.0006	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13	1.06×10^7	B	32

ACKNOWLEDGMENTS

The authors gratefully acknowledge Alexander Kramida, Atomic Spectroscopy Group, National Institute of Standards and Technology (NIST), USA for generously providing his calculations of ionization energy, LOPT code and other information. One of the authors U. D. acknowledges kind help from Arun K. Thazathveetil, Department of Chemistry, North-Western University, Evanston, USA, S. Knoop, Vrije Universiteit, Amsterdam, The Netherlands, Lotje Wansbeek, The Netherlands and B. Santra, Technische Universität Kaiserslautern, Germany.

[1] P. Curie, M. Curie, and G. Bemont, Compt. Rend. Acad. Sci. **127**, 1215 (1898).
[2] W. Ritz, Phys. Zeit. **9**, 521 (1908).
[3] David R. Lide, *CRC Hand book of Chemistry and Physics* (88^{th} Ed., CRC Press, Boca Raton, FL, 2007).
[4] E. Fues, Ann. Phys. **368**, 1 (1920).
[5] A. Fowler, *Report on Series in Line Spectra* (Fleet way press, London, 1922).
[6] E. Rasmussen, Z. Phys. **86**, 24 (1933).
[7] E. Rasmussen, Z. Phys. A **87**, 607 (1934).
[8] H. N. Russell, Phys. Rev. **46**, 989 (1934).
[9] C. E. Moore, "Atomic energy levels," (NBS Circular No 467, Washington, DC: US Govt. Printing, 1958) pp. 231–2.
[10] J. R. Guest, N. D. Scielzo, I. Ahmad, K. Bailey, J. P. Greene, R. J. Holt, Z.-T. Lu, T. P. O. 'Connor, and D. H. Potterveld, Phys. Rev. Lett. **98**, 093001 (2007).
[11] R. H. Parker, M. R. Dietrich, K. Bailey, J. P. Greene, R. J. Holt, M. R. Kalita, W. Korsch, Z.-T. Lu, P. Mueller, T. P. O. Ćonnor, J. Singh, I. A. Sulai, and W. L. Trimble, Phys. Rev. C **86**, 065503 (2012).
[12] B. Santra, U. Dammalapati, A. Groot, K. Jungmann, and L. Willmann, Phys. Rev. A **90**, 040501(R) (2014).
[13] M. Nuñez Portela, E. A. Dijck, A. Mohanty, H. Bekker, J. E. van den Berg, G. S. Giri, S. Hoekstra, C. J. G. Onderwater, O. O. Versolato, *et al.*, Appl. Phys. B **114**, 173 (2014).
[14] R. H. Parker, M. R. Dietrich, M. R. Kalita, N. D. Lemke, K. G. Bailey, M. Bishof, J. P. Greene, R. J. Holt, W. Korsch, Z.-T. Lu, P. Mueller, T. P. O. Ćonnor, and J. T. Singh, Phys. Rev. Lett. **114**, 233002 (2015).
[15] S. J. Park, Y. J. Kim, T. Shin, Y. K. Kim, and S. K. Kim, J. Phys. B: At. Mol. Opt. Phys. **47**, 245301 (2014).
[16] K. M. Lynch, J. Billowes, M. L. Bissell, I. Budinčević, T. E. Cocolios, T. D. Goodacre, R. P. de Groote, V. N. Fedosseev, K. T. Flangan, S. Franchoo, *et al.*, Proposal for collinear resonance ionization spectroscopy of radium ions INTC-P-413 (CERN, 2014).
[17] J. A. Armstrong, J. J. Wynne, and F. S. Tomkins, J. Phys. B **13**, L133 (1980).
[18] F. S. Tomkins and B. Ercoli, App. Opt. **6**, 1299 (1967).
[19] S. A. Ahmad, W. Klempt, R. Neugart, E. W. Otten, K. Wendt, C. Ekström, and The ISOLDE Collaboration, Phys. Lett. B **133**, 47 (1983).
[20] E. Arnold, W. Borchers, M. Carré, H. T. Duong, P. Juncar, J. Lermé, S. Liberman, W. Neu, R. Neugart, E. W. Otten, M. Pellarin, J. Pinard, G. Ulm, J. L. Vialle, and K. Wendt, Phys. Rev. Lett. **59**, 771 (1987).
[21] K. Wendt, S. A. Ahmad, W. Klempt, R. Neugart, E. W. Otten, and H. H. Stroke, Z. Phys. D **4**, 227 (1987).
[22] S. A. Ahmad, W. Klempt, R. Neugart, E. W. Otten, P.-G. Reinhard, G. Ulm, K. Wendt, and The ISOLDE Collaboration, Nucl. Phys. A **483**, 244 (1988).
[23] W. Neu, R. Neugart, E. W. Otten, G. Passler, K. Wendt, B. Fricke, E. Arnold, H. J. Kluge, G. Ulm, and The ISOLDE Collaboration, Z. Phys. D **11**, 105 (1989).
[24] V. V. Flambaum, Phys. Rev. A **60**, R2611 (1999).
[25] V. A. Dzuba, V. V. Flambaum, and J. S. M. Ginges, Phys. Rev. A **61**, 062509 (2000).
[26] J. Bieroń, P. Indelicato, and P. Jönsson, Eur. Phys. J. Special Topics **144**, 75 (2007).
[27] J. Bieroń, G. Gaigalas, E. Gaidamauskas, S. Fritzsche, P. Indelicato, and P. Jönsson, Phys. Rev. A **80**, 012513 (2009).
[28] J. Engel, M. Bender, J. Dobaczewski, J. H. de Jesus, and P. Olbratowski, Phys. Rev. C **68**, 025501 (2003).
[29] J. Dobaczewski and J. Engel, Phys. Rev. Lett. **94**, 232502 (2005).
[30] L. Radžiūtė, G. Gaigalas, P. Jönsson, and J. Bieroń, Phys. Rev. A **90**, 012528 (2014).
[31] L. W. Wansbeek, B. K. Sahoo, R. G. E. Timmermans, K. Jungmann, B. P. Das, and D. Mukherjee, Phys. Rev. A **78**, 050501(R) (2008).
[32] B. K. Sahoo, L. W. Wansbeek, K. Jungmann, and R. G. E. Timmermans, Phys. Rev. A **79**, 052512 (2009).
[33] O. O. Versolato, L. W. Wansbeek, K. Jungmann, R. G. E. Timmermans, L. Willmann, and H. W. Wilschut, Phys. Rev. A **83**, 043829 (2011).
[34] G. P. A. Berg, O. C. Dermois, U. Dammalapati, P. Dendooven, M. N. Harakeh, K. Jungmann, C. J. G. Onderwater, A. Rogachevskiy, M. Sohani, E. Traykov, L. Willmann, and H. Wilschut, Nucl. Instr. Meth. A **560**, 169 (2006).
[35] E. Traykov, A. Rogachevskiy, M. Bosswell, U. Dammalapati, P. Dendooven, O. C. Dermois, K. Jungmann, C. J. G. Onderwater, M. Sohani, L. Willmann, H. W. Wilschut, and A. R. Young, Nucl. Instr. Meth. B **572**, 580 (2007).
[36] P. D. Shidling, G. S. Giri, D. J. van der Hoek, K. Jungmann, W. Kruithof, C. J. G. Onderwater, M. Sohani, O. O. Versolato, L. Willmann, and H. W. Wilschut, Nucl. Instr. Meth. A **606**, 305 (2009).
[37] O. O. Versolato, G. S. Giri, L. W. Wansbeek, J. E. van den Berg,

D. J. van der Hoek, K. Jungmann, W. L. Kruithof, C. J. G. Onderwater, B. K. Sahoo, B. Santra, P. D. Shidling, R. G. E. Timmermans, L. Willman, and H. W. Wilschut, Phys. Rev. A **82**, 010501(R) (2010).
[38]O. O. Versolato, G. S. Giri, J. van den Berg, O. Böll, U. Dammalapati, D. J. van der Hoek, S. Hoekstra, K. Jungmann, W. L. Kruithof, S. Müller, M. N. Portela, C. J. G. Onderwater, B. Santra, R. G. E. Timmermans, L. W. Wansbeek, L. Willmann, and H. W. Wilschut, Phys. Lett. A **375**, 3130 (2011).
[39]G. S. Giri, O. O. Versolato, J. van de Berg, O. Böll, U. Dammalapati, D. J. van der Hoek, K. Jungmann, W. L. Kruithof, S. Müller, M. N. Portela, C. J. G. Onderwater, B. Santra, R. G. E. Timmermans, L. W. Wansbeek, L. Willmann, and H. W. Wilschut, Phys. Rev. A **84**, 020503(R) (2011).
[40]B. Santra, *Precision spectroscopy of neutral radium: Towards searches for permanent electric dipole moments*, Ph.D. thesis, University of Groningen (2012).
[41]N. D. Scielzo, J. R. Guest, E. C. Schulte, I. Ahmad, K. Bailey, D. L. Bowers, R. J. Holt, Z.-T. Lu, T. P. Oconnor, and D. H. Potterveld, Phys. Rev. A **73**, 010501(R) (2006).
[42]W. L. Trimble, I. A. Sulai, I. Ahmad, K. Bailey, B. Graner, J. P. Greene, R. J. Holt, W. Korsch, Z.-T. Lu, P. Mueller, and T. P. O. 'Connor, Phys. Rev. A **80**, 054501 (2009).
[43]E. R. Peck and K. Reeder, J. Opt. Soc. Am. **62**, 958 (1972).
[44]A. E. Kramida, Comp. Phys. Commun. **182**, 419 (2011).
[45]J. J. Curry, J. Phys. Chem. Ref. Data **33**, 725 (2004).
[46]P. J. Mohr, B. N. Taylor, and B. B. Newell, "The 2014 CODATA Recommended Values of the Fundamental Physical Constants" (Web Version 7.0). `http://physics.nist.gov/constants`, National Institute of Standards and Technology, Gaithersburg, MD 20899 (2015).
[47]J. K. Tuli, *Nuclear Wallet Cards* (Nuclear Data Center, USA, 2011).
[48]M. Wang, G. Audi, A. H. Wapstra, F. G. Kondev, M. MacCormick, X. Xu, and B. Pfeiffer, Chinese Phys. C **36**, 1603 (2012).
[49]A. Kramida, private communication.
[50]H. C. Urey, F. G. Brickwedde, and G. M. Murphy, Phys. Rev. **40**, 1 (1932).
[51]P. Quinet, C. Argante, V. Fivet, C. Terranova, A. V. Yushchenko, and È. Bièmont, A & A **474**, 307 (2007).
[52]J. S. M. Ginges and V. A. Dzuba, Phys. Rev. A **91**, 042505 (2015).
[53]V. A. Dzuba and V. V. Flambaum, J. Phys. B: At. Mol. Opt. Phys. **40**, 227 (2007).
[54]J. Bieroń, C. F. Fischer, S. Fritzsche, and K. Pachucki, J. Phys. B: At. Mol. Opt. Phys. **37**, L305 (2004).
[55]C. Sansonetti, private communication.
[56]A. Kramida, Fusion Sci. Technol. **63**, 313 (2013).
[57]V. A. Dzuba, V. V. Flambaum, and J. S. M. Ginges, Phys. Rev. A **63**, 062101 (2001).
[58]R. Pal, D. Jiang, M. S. Safronova, and U. I. Safronova, Phys. Rev. A **79**, 062505 (2009).
[59]B. K. Sahoo, B. P. Das, R. K. Chaudhuri, D. Mukherjee, R. G. Timmermans, and K. Jungmann, Phys. Rev. A **76**, 040504(R) (2007).
[60]From experiment. Accuracy, B – uncertainty is $\leq 10\%$.
[61]'p' means 'perturbed by' with the perturbing species given in parentheses.
[62]6 is the reference for the line. HQ - Hilger Quartz, Q - Quartz prism, P - Glass prism and G - Grating (used in the red region) are the letter codes of the spectrometer used. See text for details of Table X.
[63]Accuracy, A – Uncertainty is $\leq 3\%$, B – uncertainty is $\leq 10\%$, C – uncertainty is $\leq 25\%$ and D – uncertainty $\leq 54\%$.

TABLE IX. Energy levels of ^{226}Ra II.

Configuration	Term	J	Level (cm^{-1})	Uncertainty (cm^{-1})	Level LOPT (cm^{-1})	Unc. 1 LOPT (cm^{-1})	Unc. 2 LOPT (cm^{-1})	Level reference
7s	^{2}S	1/2	0.000		0.000	0.0007	0	
6d	^{2}D	3/2	12084.2721		12084.2721	0.0006	0.0010	9
	^{2}D	5/2	13742.99		13742.99	0.03	0.03	9
7p	^{2}P^0	1/2	21351.3259	0.0008	21351.3259	0.0005	0.0008	9, 13
	^{2}P^0	3/2	26208.8538	0.0021	26208.8538	0.0021	0.0021	9, 13
8s	^{2}S	1/2	43405.05		43405.05	0.05	0.05	9
7d	^{2}D	3/2	48744.04		48744.04	0.10	0.08	9
	^{2}D	5/2	49240.45		49240.45	0.18	0.4	9
5f	^{2}F^0	5/2	48987.86		48987.86	0.04	0.15	9
	^{2}F^0	7/2	49272.14		49272.14	0.04	0.16	9
8p	^{2}P^0	1/2	50606.29		50606.29	0.16	0.22	9
	^{2}P^0	3/2	52392.02		52392.02	0.10	0.19	9
9s	^{2}S	1/2	59165.3		59165.3	0.3	0.3	9
6f	^{2}F^0	5/2	59517		59517	5	5	9
	^{2}F^0	7/2	59814		59814	4	4	9
8d	^{2}D	3/2	61735.0		61735.0	0.5	0.5	9
	^{2}D	5/2	61973.8		61973.8	0.9	0.9	9
9p	^{2}P^0	3/2	63410.43		63410.43	0.20	0.21	9
5g	^{2}G	7/2,9/2	64150.49		64150.49	0.03	0.15	9
7f	^{2}F^0	5/2	66521.85		66521.85	0.22	0.24	9
	^{2}F^0	7/2	66691.18		66691.18	0.21	0.4	9
10s	^{2}S	1/2	66837.88		66838.72	0.18	0.22	9
9d	^{2}D	3/2	68264.07		68264.07	0.19	0.21	9
	^{2}D	5/2	68394.87		68394.87	0.18	0.23	9
6g	^{2}G	7/2,9/2	69560.68		69560.68	0.15	0.21	9
11s	^{2}S	1/2	71172.88		71172.88	0.21	0.3	9
10d	^{2}D	3/2	72043.33		72043.33	0.23	0.3	9
	^{2}D	5/2	72123.74		72123.74	0.19	0.3	9
7g	^{2}G	7/2,9/2	72824.21		72824.21	0.14	0.20	9
11d	^{2}D	3/2	74434.30		74434.30	0.17	0.3	9
	^{2}D	5/2	74489		74489	5	5	9
8g	^{2}G	7/2,9/2	74941.76		74941.76	0.17	0.23	9
12d	^{2}D	5/2	76080.68		76080.68	0.17	0.3	9
9g	^{2}G	7/2,9/2	76392.92		76392.92	0.15	0.21	9
10g	^{2}G	7/2,9/2	77431.2		77431.2	1.2	1.1	9
11g	^{2}G	7/2,9/2	78195.1		78195.1	2.0	1.2	9

TABLE XI. Ritz wavelengths and uncertainty of Ritz wavelengths in ^{226}Ra II

Lower level	Upper level	Ritz $\lambda/\AA$	Uncertainty Ritz $\lambda/\AA$
6d $^2D_{5/2}$	7f $^2F^0_{7/2}$	1888.639	0.015
7s $^2S_{1/2}$	8p $^2P^0_{3/2}$	1908.688	0.007
7p $^2P^0_{1/2}$	10d $^2D_{3/2}$	1972.698	0.012
7s $^2S_{1/2}$	8p $^2P^0_{1/2}$	1976.039	0.009
7p $^2P^0_{1/2}$	11s $^2S_{1/2}$	2006.514	0.011
6d $^2D_{5/2}$	9p $^2P^0_{3/2}$	2012.741	0.008
7p $^2P^0_{1/2}$	9d $^2D_{3/2}$	2130.944	0.009
6d $^2D_{5/2}$	6f $^2F^0_{7/2}$		
7p $^2P^0_{3/2}$	10d $^2D_{5/2}$	2177.261	0.013
7p $^2P^0_{3/2}$	10d $^2D_{3/2}$	2181.081	0.015
7p $^2P^0_{1/2}$	10s $^2S_{1/2}$	2197.725	0.011
7p $^2P^0_{3/2}$	11s $^2S_{1/2}$	2223.308	0.014
7p $^2P^0_{3/2}$	9d $^2D_{5/2}$	2369.730	0.013
7p $^2P^0_{3/2}$	9d $^2D_{3/2}$	2377.101	0.012
7p $^2P^0_{3/2}$	10s $^2S_{1/2}$	2460.499	0.013
6d $^2D_{3/2}$	8p $^2P^0_{3/2}$	2480.164	0.012
6d $^2D_{5/2}$	8p $^2P^0_{3/2}$	2586.613	0.013
6d $^2D_{3/2}$	8p $^2P^0_{1/2}$	2595.142	0.015
7p $^2P^0_{1/2}$	9s $^2S_{1/2}$	2643.737	0.021
6d $^2D_{3/2}$	5f $^2F^0_{5/2}$	2708.960	0.011
6d $^2D_{5/2}$	5f $^2F^0_{7/2}$	2813.762	0.012
6d $^2D_{5/2}$	5f $^2F^0_{5/2}$	2836.458	0.012
7p $^2P^0_{3/2}$	9s $^2S_{1/2}$	3033.42	0.03
5f $^2F^0_{5/2}$	11g $^2G_{7/2,9/2}$	3422.83	0.14
5f $^2F^0_{7/2}$	11g $^2G_{7/2,9/2}$	3456.47	0.14
5f $^2F^0_{5/2}$	10g $^2G_{7/2,9/2}$	3514.76	0.14
5f $^2F^0_{7/2}$	10g $^2G_{7/2,9/2}$	3550.24	0.14
7p $^2P^0_{1/2}$	7d $^2D_{3/2}$	3649.566	0.011
7s $^2S_{1/2}$	7p $^2P^0_{3/2}$	3814.4219	0.0003
5f $^2F^0_{5/2}$	8g $^2G_{7/2,9/2}$	3851.89	0.03
5f $^2F^0_{7/2}$	8g $^2G_{7/2,9/2}$	3894.55	0.03
5f $^2F^0_{5/2}$	7g $^2G_{7/2,9/2}$	4194.091	0.024
5f $^2F^0_{7/2}$	7g $^2G_{7/2,9/2}$	4244.72	0.03
7p $^2P^0_{3/2}$	7d $^2D_{5/2}$	4340.64	0.07
7p $^2P^0_{3/2}$	7d $^2D_{3/2}$	4436.259	0.017
7p $^2P^0_{1/2}$	8s $^2S_{1/2}$	4533.111	0.010
8p $^2P^0_{1/2}$	10d $^2D_{3/2}$	4663.52	0.05
7s $^2S_{1/2}$	7p $^2P^0_{1/2}$	4682.23940	0.00017
5f $^2F^0_{5/2}$	6g $^2G_{7/2,9/2}$	4859.42	0.04
5f $^2F^0_{7/2}$	6g $^2G_{7/2,9/2}$	4927.52	0.04
8s $^2S_{1/2}$	9p $^2P^0_{3/2}$	4997.26	0.05
8p $^2P^0_{3/2}$	10d $^2D_{5/2}$	5066.57	0.05
8p $^2P^0_{3/2}$	11s $^2S_{1/2}$	5323.09	0.06
8p $^2P^0_{1/2}$	9d $^2D_{3/2}$	5661.65	0.06
7d $^2D_{5/2}$	7f $^2F^0_{7/2}$	5728.83	0.07
7p $^2P^0_{3/2}$	8s $^2S_{1/2}$	5813.628	0.017
8p $^2P^0_{1/2}$	10s $^2S_{1/2}$	6158.80	0.07
8p $^2P^0_{3/2}$	9d $^2D_{5/2}$	6247.16	0.07
8p $^2P^0_{3/2}$	9d $^2D_{3/2}$	6298.64	0.06
5f $^2F_{5/2}$	5g $^2G_{7/2,9/2}$	6593.341	0.019
5f $^2F_{7/2}$	5g $^2G_{7/2,9/2}$	6719.320	0.019
8p $^2P^0_{3/2}$	10s $^2S_{1/2}$	6920.09	0.08
6d $^2D_{3/2}$	7p $^2P^0_{3/2}$	7077.9042	0.0011
6d $^2D_{5/2}$	7p $^2P^0_{3/2}$	8019.701	0.020
6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	10787.9609	0.0007

TABLE XII. Wavelengths, wavenumbers and terms of isotopes of Ra II

Atomic Mass	$\lambda_{air}/\mathring{A}$	σ/cm^{-1}	Uncertainty (cm^{-1})	Lower level	Upper level	Reference
212	3814.0962	26211.0916	0.0021	7s $^2S_{1/2}$	7p $^2P^0_{3/2}$	13
	4681.78273	21353.4085	0.0005	7s $^2S_{1/2}$	7p $^2P^0_{1/2}$	13
	7078.0714	14124.2481	0.0018	6d $^2D_{3/2}$	7p $^2P^0_{3/2}$	13
	10788.530	9266.5650	0.0004	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13
213	4681.79396	21353.3573	0.0004	7s $^2S_{1/2}$	7p $^2P^0_{1/2}$	13
	7078.0822 (F=2-F=1)	14124.2264	0.0017	6d $^2D_{3/2}$	7p $^2P^0_{3/2}$	13
	10788.5176	9266.5756	0.0005	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13
214	3814.12014	26210.9273	0.0017	7s $^2S_{1/2}$	7p $^2P^0_{3/2}$	13
	4681.81626	21353.2556	0.0005	7s $^2S_{1/2}$	7p $^2P^0_{1/2}$	13
	7078.0599	14124.2709	0.0014	6d $^2D_{3/2}$	7p $^2P^0_{3/2}$	13
	10788.4902	9266.5992	0.0004	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13
225	4682.21642	21351.4307	0.0008	7s $^2S_{1/2}$	7p $^2P^0_{1/2}$	13
	10787.9892	9267.0295	0.0006	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13
226	10787.9609	9267.0538	0.0006	6d $^2D_{3/2}$	7p $^2P^0_{1/2}$	13

TABLE XIII. Isotope shifts and hyperfine constants of 7s $^2S_{1/2}$, 7p $^2P^0_{1/2}$, and 7p $^2P^0_{3/2}$ levels in Ra II

	7s $^2S_{1/2}$	7p $^2P^0_{1/2}$		7p $^2P^0_{3/2}$			
Atomic mass	HFS A (MHz)	Isotopic frequency shift (MHz)	HFS A (MHz)	Isotopic frequency shift (MHz)[a]	HFS A (MHz)[a]	HFS B (MHz)[a]	Reference(s)
208		11950(8)					21
209							21
210		8449(6)					21
211	6624.8(1.0)	7770(4)	1299.7(0.8)				21
212		4583(3)		4937(2)			23
213	22920.0(6.0)	3049(3)	4525.0(5.0)				21
214		0		0			
220		-30808(14)					21
221	-1348.6(1.8)[a]	-36496(14)	-266.3(1.5)	-39222(18)	-22.4(9)	1364.2(5.1)	21, 23
222		-40552(14)		-43572(14)			21
223	3404.0(1.9)[a]	-45657(16)	667.1(2.1)	-49050(18)	56.5(8)	864.8(1.9)	21, 23
224		-49411(16)		-53076(21)			21
225	-27731(13)[a]	-54710(18)	-5446.0(7.0)	-58776(22)	-466.4(4.6)		21, 23
226		-57852(18)		-62159(21)			21
227	-5063.5(3.1)	-61811(19)	-996.2(2.3)				21
228		-66050(20)					21
229	3789.7(2.3)	-70432(22)	743.9(1.2)				21
230		-75456(23)					21
232							

[a] From reference 23

TABLE XIV. Isotope shifts and hyperfine structure constants A and B of the 6d $^2D_{3/2}$ and 7p $^2P^0_{1/2}$ levels in Ra II

	6d $^2D_{3/2}$		7p $^2P^0_{1/2}$		
Atomic mass	HFS A (MHz)	HFS B (MHz)	Isotopic frequency shift (MHz)	HFS A (MHz)	Reference(s)
209	148(10)[a]	104(38)[a]	2645(56)[b]	1276(20)[d]	37, 38, 39
210			1884(16)		39
211	151(2)[a]	103(6)[a]	1755(14)[b]	1299.7(0.8)[c]	38, 39, 21
212			1025(12)		39
213	528(5)[d]		707(14)[b]	4525(5)[c]	37, 39, 21
214			0		

[a] Reference 38
[b] Reference 39
[c] Reference 21
[d] Reference 37

TABLE XV. Isotope shifts and hyperfine structure constants A and B of the 7p $^2P^0_{3/2}$ and 6d $^2D_{3/2}$ levels in Ra II

	6d $^2D_{3/2}$	7p $^2P^0_{3/2}$			
Atomic mass	HFS A(MHz)	Isotopic frequency shift (MHz)	HFS A (MHz)	HFS B (MHz)	Reference(s)
212	528(5)	701(50)			37, 39
213		453(34)			37
214		0			

TABLE XVI. Lifetimes of energy levels in ^{212}Ra II

Level	6d $^2D_{5/2}$
Lifetime	232(4) ms
Reference	37